

J = 0

In the following H^0 refers to the signal that has been discovered in the Higgs searches. Whereas the observed signal is labeled as a spin 0 particle and is called a Higgs Boson, the detailed properties of H^0 and its role in the context of electroweak symmetry breaking need to be further clarified. These issues are addressed by the measurements listed below.

Concerning mass limits and cross section limits that have been obtained in the searches for neutral and charged Higgs bosons, see the sections "Searches for Neutral Higgs Bosons" and "Searches for Charged Higgs Bosons (H^{\pm} and $H^{\pm\pm}$)", respectively.

NODE=S126

HO MASS

A combination of the results by the ATLAS and CMS collaborations, which is not yet published, yields an average value of 125.09 \pm 0.21 \pm 0.11 GeV.

VALUE (GeV) DOCUMENT ID TECN COMMENT **125.13\pm0.28 OUR AVERAGE** Error includes scale factor of 1.2. [125.7 \pm 0.4 GeV OUR 2014 AVERAGE] 1,2 AAD $125.36 \pm 0.37 \pm 0.18$ 14W ATLS *pp*, 7, 8 TeV 3 CHATRCHYAN 14AA CMS pp, 7, 8 TeV, $ZZ^* \rightarrow 4\ell$ $125.6 \pm 0.4 \pm 0.2$ 4 KHACHATRY...14P CMS pp, 7, 8 TeV, $\gamma\gamma$ $124.70 \pm 0.31 \pm 0.15$

• • • We do not use the	tolle	owing data for averages, fits, limits, etc. • • •
$125.98\!\pm\!0.42\!\pm\!0.28$		1 AAD 1 14W ATLS p p , 7, 8 TeV, $^{\gamma}$ $^{\gamma}$
$124.51\!\pm\!0.52\!\pm\!0.06$		1 AAD 14W ATLS pp , 7, 8 TeV, $ZZ^* ightarrow 4\ell$
122 ± 7		⁵ CHATRCHYAN 14K CMS pp , 7, 8 TeV, $\tau\tau$
125.5 $\pm 0.2 ^{+ 0.5}_{- 0.6}$		^{2,6} AAD 13AK ATLS <i>pp</i> , 7, 8 TeV
$126.8 \pm 0.2 \pm 0.7$		6 AAD 13AK ATLS pp , 7, 8 TeV, $\gamma\gamma$
$124.3 \begin{array}{l} +0.6 & +0.5 \\ -0.5 & -0.3 \end{array}$		⁶ AAD 13AK ATLS pp , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
125.8 ± 0.4 ± 0.4		^{2,7} CHATRCHYAN 13」 CMS pp, 7, 8 TeV
$126.2 \pm 0.6 \pm 0.2$		7 CHATRCHYAN 13J CMS pp , 7, 8 TeV, $ZZ^* \rightarrow 4\ell$
$126.0 \pm 0.4 \pm 0.4$		^{2,8} AAD 12AI ATLS <i>pp</i> , 7, 8 TeV
$125.3 \pm 0.4 \pm 0.5$		^{2,9} CHATRCHYAN 12N CMS pp, 7, 8 TeV
1	1	1

 1 AAD 14W use 4.5 fb $^{-1}$ of pp collisions at $E_{\rm cm}=$ 7 TeV and 20.3 fb $^{-1}$ at 8 TeV.

²Combined value from $\gamma\gamma$ and $ZZ^* \rightarrow 4\ell$ final states.

 3 CHATRCHYAN 14AA use 5.1 fb $^{-1}$ of pp collisions at $E_{
m cm}=$ 7 TeV and 19.7 fb $^{-1}$ at $E_{\rm cm}=8$ TeV.

 4 KHACHATRYAN 14P use 5.1 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 19.7 fb $^{-1}$ at

 5 CHATRCHYAN 14K use 4.9 fb $^{-1}$ of pp collisions at $E_{\rm cm}=$ 7 TeV and 19.7 fb $^{-1}$ at 6 AAD 13AK use 4.7 fb $^{-1}$ of pp collisions at $E_{\rm cm}=$ 7 TeV and 20.7 fb $^{-1}$ at $E_{\rm cm}=$ 8 TeV.

Superseded by AAD 14W.

 7 CHATRCHYAN 13J use 5.1 fb $^{-1}$ of pp collisions at $E_{
m cm}=$ 7 TeV and 12.2 fb $^{-1}$ at $E_{
m cm}=$ 8 TeV.

 8 AAD 12AI obtain results based on 4.6–4.8 fb $^{-1}$ of pp collisions at $E_{
m cm}=$ 7 TeV and 5.8–5.9 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. An excess of events over background with a local significance of 5.9 σ is observed at $m_{H^0}=126$ GeV. See also AAD 12DA.

 9 CHATRCHYAN 12N obtain results based on 4.9–5.1 fb $^{-1}$ of $\it pp$ collisions at $\it E_{
m cm}=7$ TeV and 5.1–5.3 fb $^{-1}$ at $E_{\rm cm}=$ 8 TeV. An excess of events over background with a local significance of 5.0 σ is observed at about $m_{\mbox{$H^0$}}=$ 125 GeV. See also CHATRCHYAN 12BY and CHATRCHYAN 13Y.

NODE=S126M NODE=S126M

NODE=S126M NEW

OCCUR=2

OCCUR=3

OCCUR=2 OCCUR=3

OCCUR=2

NODE=S126M;LINKAGE=A NODE=S126M;LINKAGE=AA NODE=S126M:LINKAGE=B

NODE=S126M;LINKAGE=C

NODE=S126M;LINKAGE=D

NODE=\$126M;LINKAGE=LH

NODE=S126M;LINKAGE=CT

NODE=S126M;LINKAGE=AI

NODE=S126M:LINKAGE=CH

HO SPIN AND CP PROPERTIES

The observation of the signal in the $\gamma\gamma$ final state rules out the possibility that the discovered particle has spin 1, as a consequence of the Landau-Yang theorem. This argument relies on the assumptions that the decaying particle is an on-shell resonance and that the decay products are indeed two photons rather than two pairs of boosted photons, which each could in principle be misidentified as a single photon.

NODE=S126CP NODE=S126CP Concerning distinguishing the spin 0 hypothesis from a spin 2 hypothesis, some care has to be taken in modelling the latter in order to ensure that the discriminating power is actually based on the spin properties rather than on unphysical behavior that may affect the model of the spin 2 state.

Under the assumption that the observed signal consists of a single state rather than an overlap of more than one resonance, it is sufficient to discriminate between distinct hypotheses in the spin analyses. On the other hand, the determination of the *CP* properties is in general much more difficult since in principle the observed state could consist of any admixture of *CP*-even and *CP*-odd components. As a first step, the compatibility of the data with distinct hypotheses of pure *CP*-even and pure *CP*-odd states with different spin assignments has been investigated. In order to treat the case of a possible mixing of different *CP* states, certain cross section ratios are considered. Those cross section ratios need to be distinguished from the amount of mixing between a *CP*-even and a *CP*-odd state, as the cross section ratios depend in addition also on the coupling strengths of the *CP*-even and *CP*-odd components to the involved particles. A small relative coupling implies a small sensitivity of the corresponding cross section ratio to effects of *CP* mixing.

VALUE <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

 \bullet \bullet We do not use the following data for averages, fits, limits, etc. \bullet \bullet

 1 ABAZOV 14F compare the $J^{CP}=0^+$ Standard Model assignment with $J^{CP}=0^-$ and 2^+ (graviton-like coupling) hypotheses in up to 9.7 fb $^{-1}$ of $p\bar{p}$ collisions at $E_{\rm Cm}=1.96$ TeV. They use kinematic correlations between the decay products of the vector boson and the Higgs boson in the final states $ZH\to\ell\ell b\bar{b},~WH\to\ell\nu b\bar{b},$ and $ZH\to\nu\nu b\bar{b}.$ The 0^- (2 $^+$) hypothesis is excluded at 97.6% CL (99.0% CL). In order to treat the case of a possible mixture of a 0^+ state with another J^{CP} state, the cross section fractions $f_X=\sigma_X/(\sigma_{0^+}+\sigma_X)$ are considered, where $X=0^-,~2^+.$ Values for f_{0^-} (f_{2^+}) above 0.80 (0.67) are excluded at 95% CL.

² CHATRCHYAN 14AA compare the $J^{CP}=0^+$ Standard Model assignment with various J^{CP} hypotheses in 5.1 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 19.7 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. $J^{CP}=0^-$ and 1^\pm hypotheses are excluded at 99% CL, and several J=2 hypotheses are excluded at 95% CL. In order to treat the case of a possible mixture of a 0^+ state with another J^{CP} state, the cross section fraction $f_{a3}=|a_3|^2 \sigma_3 \ / \ (|a_1|^2 \sigma_1 + |a_2|^2 \sigma_2 + |a_3|^2 \sigma_3)$ is considered, where the case $a_3=1$, $a_1=a_2=0$ corresponds to a pure CP-odd state. Assuming $a_2=0$, a value for f_{a3} above 0.51 is excluded at 95% CL.

^{95/6} CL. ³ CHATRCHYAN 14G compare the $J^{CP}=0^+$ Standard Model assignment with $J^{CP}=0^-$ and 2^+ (graviton-like coupling) hypotheses in 4.9 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 19.4 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. Varying the fraction of the production of the 2^+ state via gg and $q\bar{q}$, 2^+ hypotheses are disfavored at CL between 83.7 and 99.8%. The 0^- hypothesis is disfavored against 0^+ at the 65.3% CL.

⁴ KHACHATRYAN 14P compare the $J^{CP}=0^+$ Standard Model assignment with a 2^+ (graviton-like coupling) hypothesis in 5.1 fb⁻¹ of pp collisions at $E_{\rm cm}=7$ TeV and 19.7 fb⁻¹ at $E_{\rm cm}=8$ TeV. Varying the fraction of the production of the 2^+ state via gg and $q\overline{q}$, 2^+ hypotheses are disfavored at CL between 71 and 94%.

⁵ AAD 13AJ compare the spin 0, *CP*-even hypothesis with specific alternative hypotheses of spin 0, *CP*-odd, spin 1, *CP*-even and *CP*-odd, and spin 2, *CP*-even models using the Higgs boson decays $H \to \gamma \gamma$, $H \to ZZ^* \to 4\ell$ and $H \to WW^* \to \ell\nu\ell\nu$ and combinations thereof. The data are compatible with the spin 0, *CP*-even hypothesis, while all other tested hypotheses are excluded at confidence levels above 97.8%.

⁶ CHATRCHYAN 13J study angular distributions of the lepton pairs in the ZZ^* channel where both Z bosons decay to e or μ pairs. Under the assumption that the observed particle has spin 0, the data are found to be consistent with the pure CP-even hypothesis, while the pure CP-odd hypothesis is disfavored.

NODE=S126CP

NODE=S126CP;LINKAGE=AB

NODE=S126CP;LINKAGE=A

NODE=S126CP;LINKAGE=C

NODE=S126CP;LINKAGE=B

NODE=S126CP;LINKAGE=AA

NODE=S126CP;LINKAGE=CH

HO DECAY WIDTH

The total decay width for a light Higgs boson with a mass in the observed range is not expected to be directly observable at the LHC. For the case of the Standard Model the prediction for the total width is about 4 MeV, which is three orders of magnitude smaller than the experimental mass resolution. There is no indication from the results

NODE=S126W NODE=S126W observed so far that the natural width is broadened by new physics effects to such an extent that it could be directly observable. Furthermore, as all LHC Higgs channels rely on the identification of Higgs decay products, the total Higgs width cannot be measured indirectly without additional assumptions. The different dependence of on-peak and off-peak contributions on the total width in Higgs decays to ZZ^* and interference effects between signal and background in Higgs decays to $\gamma\gamma$ can provide additional information in this context. Constraints on the total width from the combination of on-peak and off-peak contributions in Higgs decays to ZZ^* rely on the assumption of equal on- and off-shell effective couplings. Without an experimental determination of the total width or further theoretical assumptions, only ratios of couplings can be determined at the LHC rather than absolute values of couplings.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
< 5.0	95	¹ AAD 14W	ATLS	pp, 7, 8 TeV, $\gamma\gamma$
< 2.6	95			pp , 7, 8 TeV, $ZZ^* ightarrow 4\ell$
<3.4	95			pp , 7, 8 TeV, $ZZ^* ightarrow 4\ell$
<2.4	95	³ KHACHATRY14P	CMS	$ ho ho$, 7, 8 TeV, $\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁴ KHACHATRY...14D CMS pp, 7, 8 TeV, $ZZ^{(*)}$

 1 AAD 14W use 4.5 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 20.3 fb $^{-1}$ at 8 TeV. The expected limit is 6.2 GeV.

 2 CHATRCHYAN 14AA use 5.1 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 19.7 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The expected limit is 2.8 GeV.

 3 KHACHATRYAN 14P use 5.1 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 19.7 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The expected limit is 3.1 GeV.

 4 KHACHATRYAN 14D derive constraints on the total width from comparing $ZZ^{(st)}$ production via on-shell and off-shell H^0 . 4ℓ and $\ell\ell\nu\nu$ final states in 5.1 fb⁻¹ of pp collisions at $E_{\rm cm}=7~{\rm TeV}$ and $19.7~{\rm fb}^{-1}$ at $E_{\rm cm}=8~{\rm TeV}$ are used.

NODE=S126W

OCCUR=2

NODE=S126W;LINKAGE=AA

NODE=S126W;LINKAGE=B

NODE=S126W;LINKAGE=C

NODE=S126W;LINKAGE=A

H⁰ DECAY MODES

	Mode	Fraction (Γ_i/Γ)	Confidence level	
$\overline{\Gamma_1}$	WW*			DESIG=1
Γ_2^-	<i>Z Z</i> *			DESIG=2
Γ_3	$\gamma \gamma$			DESIG=3
Γ ₄	$b\overline{b}$			DESIG=4
Γ_5	$\mu^+\mu^-$			DESIG=8
Γ_6	$ au^+ au^-$			DESIG=5
Γ_7	$Z\gamma$			DESIG=6
Γ ₈	invisible	<58 %	95%	DESIG=7

NODE=S126220;NODE=S126

 Γ_8/Γ

H⁰ BRANCHING RATIOS

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$

Invisible final states. TECN COMMENT

140 ATLS $pp \rightarrow H^0ZX$, 7, 8 TeV < 0.75 95 2 CHATRCHYAN 14B CMS $pp \rightarrow H^{0}ZX$, $qqH^{0}X$ < 0.58 95

ullet ullet We do not use the following data for averages, fits, limits, etc. ullet ullet

 3 CHATRCHYAN14B CMS $pp \rightarrow H^0ZX$, 7, 8 TeV 4 CHATRCHYAN14B CMS $pp \rightarrow qqH^0X$, 8 TeV < 0.81 < 0.65

¹ AAD 140 search for $pp \to H^0 ZX$, $Z \to \ell \ell$, with H^0 decaying to invisible final states in 4.5 fb $^{-1}$ at $E_{\rm cm}=$ 7 TeV and 20.3 fb $^{-1}$ at $E_{\rm cm}=$ 8 TeV. The quoted limit on the branching ratio is given for $m_{H^0}=$ 125.5 GeV and assumes the Standard Model rate for

² CHATRCHYAN 14B search for $pp \to H^0 ZX$, $Z \to \ell\ell$ and $Z \to b\overline{b}$, and also $pp \to qqH^0 X$ with H^0 decaying to invisible final states using data at $E_{\rm cm}=7$ and 8 TeV. The quoted limit on the branching ratio is obtained from a combination of the limits from $H^0 Z$ and $q q H^0$. It is given for $m_{H^0} = 125~{\rm GeV}$ and assumes the Standard Model rates for the two production processes.

³ CHATRCHYAN 14B search for $pp \to H^0 ZX$ with H^0 decaying to invisible final states and $Z \to \ell \ell$ in 4.9 fb⁻¹ at $E_{\rm cm} = 7$ TeV and 19.7 fb⁻¹ at $E_{\rm cm} = 8$ TeV, and also with $Z \to b \overline{b}$ in 18.9 fb⁻¹ at $E_{\rm cm} = 8$ TeV. The quoted limit on the branching ratio is given for $m_{H^0}=125~{\rm GeV}$ and assumes the Standard Model rate for $H^0\,Z$ production.

NODE=S126225

NODE=S126R01

NODE=S126R01 NODE=S126R01;CHECK LIMITS

OCCUR=2 OCCUR=3

NODE=S126R01;LINKAGE=A

NODE=S126R01;LINKAGE=B

NODE=S126R01;LINKAGE=CH

 4 CHATRCHYAN 14B search for $p\,p\to q\,q\,H^0\,X$ (vector boson fusion) with H^0 decaying to invisible final states in 19.5 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The quoted limit on the branching ratio is given for $m_{H^0}=125$ GeV and assumes the Standard Model rate for $q\,q\,H^0$ production.

NODE=S126R01;LINKAGE=C

HO SIGNAL STRENGTHS IN DIFFERENT CHANNELS

The H^0 signal strength in a particular final state xx is given by the cross section times branching ratio in this channel normalized to the Standard Model (SM) value, $\sigma \cdot \mathsf{B}(H^0 \to xx) / (\sigma \cdot \mathsf{B}(H^0 \to xx))_{\text{SM}}$, for the specified mass value of H^0 .

NODE=S126230

NODE=S126230

Combined Final States

VALUE	DOCUMENT ID	TECN	COMMENT
1.17 ± 0.17 OUR AVERAGE	Error includes scale fa	ctor of 1.2.	
$1.33^{\begin{subarray}{c} +0.14 \\ -0.10 \end{subarray}} \pm 0.15$	¹ AAD	13AK ATLS	<i>pp</i> , 7 and 8 TeV
$1.44 ^{igoplus 0.59}_{-0.56}$	² AALTONEN	13M TEVA	$p\overline{p} \rightarrow H^0X$, 1.96 TeV
0.87 ± 0.23	³ CHATRCHYAI	N 12N CMS	$pp \rightarrow H^0 X$, 7, 8 TeV
• • • We do not use the follo	wing data for averages	s, fits, limits, e	etc. • • •

NODE=S126SA NODE=S126SA

OCCUR=2 OCCUR=3

 1 AAD 13AK use 4.7 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 20.7 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The combined signal strength is based on the $\gamma\gamma,~ZZ^*\to~4\ell,$ and $WW^*\to\ell\nu\ell\nu$ channels. The quoted signal strength is given for $m_{\mbox{\it H}^0}=125.5$ GeV. Reported statistical error value modified following private communication with the experiment.

NODE=S126SA;LINKAGE=LH

² AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to 10.0 fb⁻¹ and 9.7 fb⁻¹, respectively, of $p\overline{p}$ collisions at $E_{\rm cm}=1.96$ TeV. The quoted signal strength is given for $m_{H^0}=125$ GeV.

 ${\sf NODE=S126SA;LINKAGE=AT}$

 3 CHATRCHYAN 12N obtain results based on 4.9–5.1 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 5.1–5.3 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. An excess of events over background with a local significance of 5.0 σ is observed at about $m_{H^0}=125$ GeV. The combined signal strength is based on the $\gamma\gamma$, ZZ^* , WW^* , $\tau^+\tau^-$, and $b\overline{b}$ channels. The quoted signal strength is given for $m_{H^0}=125.5$ GeV. See also CHATRCHYAN 13Y.

NODE=S126SA;LINKAGE=CA

 4 AALTONEN 13L combine all CDF results with 9.45–10.0 fb $^{-1}$ of $p\bar{p}$ collisions at $E_{\rm cm}=1.96$ TeV. The quoted signal strength is given for $m_{\mbox{\it H}^0}=125$ GeV.

NODE=S126SA;LINKAGE=LL

 5 ABAZOV 13L combine all D0 results with up to 9.7 fb $^{-1}$ of $p\overline{p}$ collisions at $E_{\rm cm}=1.96$ TeV. The quoted signal strength is given for $m_{H^0}=125$ GeV.

NODE=S126SA;LINKAGE=AB

 6 AAD 12AI obtain results based on 4.6–4.8 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 5.8–5.9 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. An excess of events over background with a local significance of 5.9 σ is observed at $m_{H^0}=126$ GeV. The quoted signal strengths are given for $m_{H^0}=126$ GeV. See also AAD 12DA.

NODE=\$126\$A;LINKAGE=AA

WW* Final State

VALUE	DOCUMENT ID	<u> TECN</u>	COMMENT	
0.81±0.16 OUR AVERAGE [0.87 ⁺ 0.24 OUR 2014 AVERAGE]				
$0.72\!\pm\!0.12\!\pm\!0.10\!+\!0.12\\-0.10$	¹ CHATRCHYAI	N 14G CMS	pp, 7, 8 TeV	ı
$0.99^{+0.31}_{-0.28}$	² AAD	13AK ATLS	<i>pp</i> , 7 and 8 TeV	

³ AALTONEN 13M TEVA $p\overline{p} \rightarrow H^0 X$, 1.96 TeV

NODE=S126SWW NODE=S126SWW NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.00^{+1.78}_{-0.00}$	⁴ AALTONEN	13L CDF	$p\overline{p} \to H^0X$, 1.96 TeV
$1.90 {}^{+ 1.63}_{- 1.52}$	⁵ ABAZOV	13L D0	$p\overline{p} \to H^0X$, 1.96 TeV
1.3 ±0.5	⁶ AAD	12AI ATLS	$pp \rightarrow H^0 X$, 7, 8 TeV
$0.5\ \pm0.6$	⁶ AAD	12AI ATLS	$pp \rightarrow H^0 X$, 7 TeV
1.9 ± 0.7	⁶ AAD	12AI ATLS	$pp \rightarrow H^0 X$, 8 TeV
$0.60^{+0.42}_{-0.37}$	⁷ CHATRCHYAI	N 12N CMS	$pp \rightarrow H^0 X$, 7, 8 TeV

OCCUR=2 OCCUR=3

- $^1\mathrm{CHATRCHYAN}$ 14G use 4.9 fb $^{-1}$ of pp collisions at $E_\mathrm{cm}=7$ TeV and 19.4 fb $^{-1}$ at $E_\mathrm{cm}=8$ TeV. The last uncertainty in the measurement is theory systematics. The quoted signal strength is given for $m_{H^0}=125.6$ GeV.
- 2 AAD 13AK use 4.7 fb $^{-1}$ of pp collisions at $E_{\rm cm}=$ 7 TeV and 20.7 fb $^{-1}$ at $E_{\rm cm}=$ 8 TeV. The quoted signal strength is given for $m_{H^0}=$ 125.5 GeV.
- 3 AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to 10.0 fb $^{-1}$ and 9.7 fb $^{-1}$, respectively, of $p\overline{p}$ collisions at $E_{\rm cm}=1.96$ TeV. The quoted signal strength is given for $m_{\mbox{\scriptsize H^0}}=125$ GeV.
- 4 AALTONEN 13L combine all CDF results with 9.45–10.0 fb $^{-1}$ of $p\bar{p}$ collisions at $E_{\rm cm}=1.96$ TeV. The quoted signal strength is given for $m_{\mbox{\it H}^0}=125$ GeV.
- 5 ABAZOV 13L combine all D0 results with up to 9.7 fb $^{-1}$ of $p\bar{p}$ collisions at $E_{\rm cm}=1.96$ TeV. The quoted signal strength is given for $m_{H^0}=125$ GeV.
- 6 AAD 12AI obtain results based on 4.7 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 5.8 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The quoted signal strengths are given for $m_{\mbox{$H^0$}}=126$ GeV. See _also AAD 12DA.
- ⁷ CHATRCHYAN 12N obtain results based on 4.9 fb⁻¹ of pp collisions at $E_{\rm cm}=7$ TeV and 5.1 fb⁻¹ at $E_{\rm cm}=8$ TeV. The quoted signal strength is given for $m_{H^0}=125.5$ GeV. See also CHATRCHYAN 13Y.

ZZ* Final State

 \bullet \bullet We do not use the following data for averages, fits, limits, etc. \bullet \bullet

	³ AAD	14AR ATLS	<i>pp</i> , 8 TeV
$1.43^{+0.40}_{-0.35}$	⁴ AAD	13AK ATLS	pp, 7 and 8 TeV
$0.80^{igoplus 0.35}_{igoplus 0.28}$	⁵ CHATRCHYAN	N 13J CMS	$pp \rightarrow H^0 X$, 7, 8 TeV
1.2 ± 0.6	⁶ AAD	12AI ATLS	$pp \rightarrow H^0 X$, 7, 8 TeV
1.4 ± 1.1	⁶ AAD	12AI ATLS	$pp \rightarrow H^0 X$, 7 TeV
1.1 ± 0.8	⁶ AAD	12AI ATLS	$pp \rightarrow H^0 X$, 8 TeV
$0.73^{+0.45}_{-0.33}$	⁷ CHATRCHYAN	N12N CMS	$pp \rightarrow H^0 X$, 7, 8 TeV

- 1 AAD 15F use 4.5 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 20.3 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The quoted signal strength is given for $m_{H^0}=125.36$ GeV. The signal strength for the gluon fusion production mode is $1.66^{+0.45}_{-0.41}^{+0.25}_{-0.15}$, while the signal strength for the vector boson fusion production mode is $0.26^{+1.60}_{-0.91}^{+0.02}_{-0.23}$.
- 2 CHATRCHYAN 14AA use $5.1~{\rm fb}^{-1}$ of pp collisions at $E_{\rm cm}=7~{\rm TeV}$ and $19.7~{\rm fb}^{-1}$ at $E_{\rm cm}=8~{\rm TeV}$. The quoted signal strength is given for $m_{H^0}=125.6~{\rm GeV}$. The signal strength for the gluon fusion and $t\bar{t}H$ production mode is $0.80^{+0.46}_{-0.36}$, while the signal strength for the vector boson fusion and WH^0 , ZH^0 production mode is $1.7^{+2.2}_{-2.1}$.
- 3 AAD 14AR measure the cross section for $pp\to H^0X,\,H^0\to ZZ^*$ using 20.3 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. They give $\sigma\cdot B=2.11^{+0.53}_{-0.47}\pm 0.08$ fb in their fiducial region, where 1.30 ± 0.13 fb is expected in the Standard Model for $m_{\slashed{H^0}}=125.4$ GeV. Various differential cross sections are also given, which are in agreement with the Standard Model expectations.
- ⁴ AAD 13AK use 4.7 fb⁻¹ of pp collisions at $E_{\rm cm}=7$ TeV and 20.7 fb⁻¹ at $E_{\rm cm}=8$ TeV. The quoted signal strength is given for $m_{H^0}=125.5$ GeV.
- 5 CHATRCHYAN 13J obtain results based on $ZZ\to 4\ell$ final states in 5.1 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 12.2 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The quoted signal strength is given for $m_{H^0}=125.8$ GeV. Superseded by CHATRCHYAN 14AA.
- 6 AAD 12AI obtain results based on 4.7–4.8 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 5.8 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The quoted signal strengths are given for $m_{H^0}=126$ GeV. See also AAD 12DA.
- 7 CHATRCHYAN 12N obtain results based on 4.9–5.1 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 5.1–5.3 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. An excess of events over background with a local significance of 5.0 σ is observed at about $m_{H^0}=125$ GeV. The quoted signal strengths are given for $m_{H^0}=125.5$ GeV. See also CHATRCHYAN 12BY and CHATRCHYAN 13Y.

NODE=S126SWW;LINKAGE=A

NODE=S126SWW;LINKAGE=LH

NODE=S126SWW;LINKAGE=AT

NODE=S126SWW;LINKAGE=LL

NODE=S126SWW;LINKAGE=AB

NODE=S126SWW;LINKAGE=AA

NODE=S126SWW;LINKAGE=CA

NODE=S126SZZ NODE=S126SZZ

NEW

OCCUR=2 OCCUR=3

NODE=S126SZZ;LINKAGE=B

NODE=S126SZZ;LINKAGE=A

NODE=S126SZZ;LINKAGE=C

NODE=S126SZZ;LINKAGE=LH

NODE=S126SZZ;LINKAGE=CA

NODE=S126SZZ;LINKAGE=AA

NODE=S126SZZ;LINKAGE=CH

7/20/2015 14:17 $\gamma\gamma$ Final State NODE=S126SGG NODE=S126SGG VALUE DOCUMENT ID TECN COMMENT $1.17^{+0.19}_{-0.17}$ OUR AVERAGE NEW $[1.58^{+0.27}_{-0.23} \ { m OUR} \ 2014 \ { m AVERAGE}]$ $1.17\!\pm\!0.23^{\,+\,0.10\,+\,0.12}_{\,-\,0.08\,-\,0.08}$ 14BC ATLS $pp \rightarrow H^0 X$, 7, 8 TeV $1.14 \!\pm\! 0.21 \!+\! 0.09 \!+\! 0.13 \\ -0.05 \!-\! 0.09$ pp, 7, 8 TeV ² KHACHATRY...14P CMS $5.97^{+3.39}_{-3.12}$ 13M TEVA $p\overline{p} \rightarrow H^0 X$, 1.96 TeV ³ AALTONEN • We do not use the following data for averages, fits, limits, etc. • • $1.55^{+0.33}_{-0.28}$ ⁴ AAD 13AK ATLS pp, 7 and 8 TeV $p\overline{p} \rightarrow H^0 X$, 1.96 TeV ⁵ AALTONEN 13L CDF $4.20^{+4.60}_{-4.20}$ ⁶ ABAZOV 13L D0 $p\overline{p} \rightarrow H^0 X$, 1.96 TeV 7 AAD 12AI ATLS $pp \rightarrow H^0 X$, 7, 8 TeV 1.8 ± 0.5 12AI ATLS $pp \rightarrow H^0 X$, 7 TeV 7 AAD 2.2 ± 0.7 OCCUR=2 7 AAD 12AI ATLS $pp \rightarrow H^0 X$, 8 TeV OCCUR=3 1.5 ± 0.6 $1.54^{+0.46}_{-0.42}$ ⁸ CHATRCHYAN 12N CMS $pp \rightarrow H^0 X$, 7, 8 TeV 1 AAD 14BC use 4.5 fb $^{-1}$ of pp collisions at $E_{\rm CM}=7$ TeV and 20.3 fb $^{-1}$ at $E_{\rm CM}=8$ TeV. The last uncertainty in the measurement is theory systematics. The quoted signal strength is given for $m_{H^0}=125.4$ GeV. The signal strengths for the individual production modes are: 1.32 ± 0.38 for gluon fusion, 0.8 ± 0.7 for vector boson fusion, 1.0 ± 1.6 for WH^0 production, $0.1_{-0.1}^{+3.7}$ for ZH^0 production, and $1.6_{-1.8}^{+2.7}$ for $t\bar{t}H^0$ resolution. NODE=S126SGG;LINKAGE=B ² KHACHATRYAN 14P use 5.1 fb⁻¹ of pp collisions at $E_{\rm cm}=7$ TeV and 19.7 fb⁻¹ at $E_{\rm cm}=8$ TeV. The last uncertainty in the measurement is theory systematics. The quoted signal strength is given for $m_{H^0}=124.7$ GeV. The signal strength for the gluon NODE=S126SGG;LINKAGE=A fusion and $t\bar{t}H$ production mode is $1.13^{+0.37}_{-0.31}$, while the signal strength for the vector boson fusion and WH^0 , ZH^0 production mode is $1.16^{+0.63}_{-0.58}$ ³ AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to 10.0 fb⁻¹ and 9.7 fb⁻¹, respectively, of $p\overline{p}$ collisions at $E_{\rm cm}=1.96$ TeV. The NODE=S126SGG;LINKAGE=AT quoted signal strength is given for $m_{H^0}=125~{\rm GeV}.$ 4 AAD 13AK use 4.7 fb $^{-1}$ of pp collisions at $E_{\rm cm}=$ 7 TeV and 20.7 fb $^{-1}$ at $E_{\rm cm}=$ 8 TeV. The quoted signal strength is given for $m_{H^0}=$ 125.5 GeV. NODE=S126SGG;LINKAGE=LH 5 AALTONEN 13L combine all CDF results with 9.45–10.0 fb $^{-1}$ of $p \overline{p}$ collisions at $E_{ m cm}$ NODE=S126SGG;LINKAGE=LL = 1.96 TeV. The quoted signal strength is given for $m_{H^0} = 125$ GeV. 6 ABAZOV 13L combine all D0 results with up to 9.7 fb $^{-1}$ of $p\overline{p}$ collisions at $E_{\rm cm}=$ NODE=S126SGG:LINKAGE=AB 1.96 TeV. The quoted signal strength is given for $m_{H^0}=125~{\rm GeV}.$ 7 AAD 12AI obtain results based on 4.8 fb $^{-1}$ of $\it pp$ collisions at $\it E_{\sf cm}=$ 7 TeV and 5.9 NODE=S126SGG;LINKAGE=AA fb $^{-1}$ at $E_{ m cm}=$ 8 TeV. The quoted signal strengths are given for $m_{H^0}=$ 126 GeV. See also AAD 12DA. 8 CHATRCHYAN 12N obtain results based on 5.1 fb $^{-1}$ of pp collisions at $E_{\rm cm}{=}7$ TeV NODE=S126SGG;LINKAGE=CA and 5.3 fb⁻¹ at $E_{\rm cm}$ =8 TeV. The quoted signal strength is given for m_{H^0} =125.5 GeV. See also CHATRCHYAN 13Y. **b** Final State NODE=S126SBB NODE=S126SBB NEW

VALUE DOCUMENT ID		<u>TECN</u>	<u>COMME</u>	NT	_	
0.85±0.29 OUR AVER						
$[1.1\pm0.5~{ m OUR}~2014~{ m A}]$	WERAGEJ					_
$0.52\!\pm\!0.32\!\pm\!0.24$	¹ AAD	15 G	ATLS	$pp \rightarrow$	H^0W/ZX , 7, 8 TeV	I
$1.0\ \pm0.5$	² CHATRCHYAN	14 AI	CMS	$pp \rightarrow$	H^0W/ZX , 7, 8 TeV	ı
$1.59^{+0.69}_{-0.72}$	³ AALTONEN	13M	TEVA	$p\overline{p} \to$	<i>H</i> ⁰ <i>X</i> , 1.96 TeV	
ullet $ullet$ We do not use the	ne following data f	or ave	erages, f	its, limit	ts, etc. • • •	
$1.72^{+0.92}_{-0.87}$	⁴ AALTONEN	13L	CDF	$p\overline{p}$ \rightarrow	<i>H</i> ⁰ <i>X</i> , 1.96 TeV	
$1.23^{+1.24}_{-1.17}$	⁵ ABAZOV	13L	D0	$p\overline{p} \to$	<i>H</i> ⁰ <i>X</i> , 1.96 TeV	
0.5 ±2.2	⁶ AAD	12AI	ATLS	$pp \rightarrow$	H^0W/ZX , 7 TeV	
	⁷ AALTONEN	12T	TEVA	$p\overline{p} \rightarrow$	$H^0 W/ZX$, 1.96 TeV	
$0.48^{+0.81}_{-0.70}$	⁸ CHATRCHYAN	12N	CMS	$pp \rightarrow$	H^0W/ZX , 7, 8 TeV	

NODE=S126STT;LINKAGE=CA

```
^1 AAD 15G use 4.7 fb^{-1} of pp collisions at E_{\rm cm}= 7 TeV and 20.3 fb^{-1} at E_{\rm cm}= 8 TeV. The quoted signal strength is given for m_{H^0}= 125.36 GeV.
                                                                                                                                 NODE=$126$BB;LINKAGE=B
   ^2 CHATRCHYAN 14AI use up to 5.1 fb^{-1} of \it pp collisions at \it E_{\rm cm}= 7 TeV and up to
                                                                                                                                 NODE=S126SBB;LINKAGE=A
     18.9 fb<sup>-1</sup> at E_{\rm cm}=8 TeV. The quoted signal strength is given for m_{H^0}=125 GeV.
     See also CHATRCHYAN 14AJ.
   AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to 10.0 fb<sup>-1</sup> and 9.7 fb<sup>-1</sup>, respectively, of p\overline{p} collisions at E_{\rm cm}=1.96 TeV. The
                                                                                                                                 NODE=S126SBB;LINKAGE=AT
     quoted signal strength is given for m_{H^0}=125 GeV.
   ^4 AALTONEN 13L combine all CDF results with 9.45–10.0 fb^{-1} of p\overline{p} collisions at E_{\rm cm}=1.96 TeV. The quoted signal strength is given for m_{\mbox{\it H}^0}=125 GeV.
                                                                                                                                 NODE=S126SBB:LINKAGE=LL
   ^{5} ABAZOV 13L combine all D0 results with up to 9.7 fb^{-1} of p\overline{p} collisions at E_{
m cm}=
                                                                                                                                 NODE=S126SBB;LINKAGE=AB
     1.96 TeV. The quoted signal strength is given for m_{H^0}=125~{\rm GeV}.
   ^6 AAD 12AI obtain results based on 4.6–4.8 fb^{-1} of pp collisions at E_{\rm cm}= 7 TeV. The quoted signal strengths are given in their Fig. 10 for m_{\mbox{\it H}^0}= 126 GeV. See also Fig. 13
                                                                                                                                 NODE=S126SBB;LINKAGE=AA
   of AAD 12DA.

AALTONEN 12T combine AALTONEN 12Q, AALTONEN 12R, AALTONEN 12s,
                                                                                                                                 NODE=S126SBB;LINKAGE=AL
     ABAZOV 120, ABAZOV 12P, and ABAZOV 12K. An excess of events over background
     is observed which is most significant in the region m_{H^0}=120–135 GeV, with a local
     significance of up to 3.3 \sigma. The local significance at m_{H^0}=125 GeV is 2.8 \sigma, which
     corresponds to (\sigma(H^0W) + \sigma(H^0Z)) \cdot B(H^0 \rightarrow b\overline{b}) = (0.23^{+0.09}_{-0.08}) pb, compared to the Standard Model expectation at m_{H^0} = 125 GeV of 0.12 \pm 0.01 pb. Superseded by
     AALTONEN 13M.
   ^{8} CHATRCHYAN 12N obtain results based on 5.0 fb^{-1} of pp collisions at E_{\rm cm}{=}7 TeV
                                                                                                                                 NODE=S126SBB;LINKAGE=CA
     and 5.1 fb<sup>-1</sup> at E_{\rm cm}=8 TeV. The quoted signal strength is given for m_{H^0}=125.5 GeV.
     See also CHATRCHYAN 13Y.
\mu^+\mu^- Final State
                                                                                                                                 NODE=S126SMU
NODE=S126SMU
                                               DOCUMENT ID TECN COMMENT
VALUE
                               CL%
ullet ullet We do not use the following data for averages, fits, limits, etc. ullet ullet
                                             <sup>1</sup> AAD
                                                                    14AS ATLS pp \rightarrow H^0 X, 7, 8 TeV
   ^1 AAD 14AS search for H^0 
ightarrow \ \mu^+\mu^- in 4.5 fb^{-1} of pp collisions at E_{
m cm}= 7 TeV and
                                                                                                                                 NODE=S126SMU;LINKAGE=A
     20.3 fb<sup>-1</sup> at E_{\rm cm}=8 TeV. The quoted signal strength is given for m_{H^0}=125.5 GeV.
	au^+	au^- Final State
                                                                                                                                 NODE=S126STT
                                                                                                                                 NODE=S126STT
VALUE
                                                DOCUMENT ID
                                                                        TECN COMMENT
0.79 ± 0.26 OUR AVERAGE
                                                                                                                                 NEW
[0.4 \pm 0.6 \text{ OUR } 2014 \text{ AVERAGE}]
                                             <sup>1</sup> CHATRCHYAN 14K CMS pp \rightarrow H^0 X, 7, 8 TeV
1.68^{igoplus 2.28}_{-1.68}
                                             <sup>2</sup> AALTONEN 13M TEVA p\overline{p} \rightarrow H^0 X, 1.96 TeV
                                             3 AAD
                                                                    12AI ATLS pp \rightarrow H^0 X, 7 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •
0.00 ^{+\, 8.44}_{-\, 0.00}
                                                                    13L CDF p\overline{p} \rightarrow H^0 X, 1.96 TeV
                                             <sup>4</sup> AALTONEN
3.96^{igoplus 4.11}_{-3.38}
                                                                                     p\overline{p} \rightarrow H^0 X, 1.96 TeV
                                                                    13L D0
0.09^{+0.76}_{-0.74}
                                             <sup>6</sup> CHATRCHYAN 12N CMS pp \rightarrow H^0 X, 7, 8 TeV
   ^1 CHATRCHYAN 14K use 4.9 fb^{-1} of pp collisions at E_{\rm cm}=7 TeV and 19.7 fb^{-1} at E_{\rm cm}=8 TeV. The quoted signal strength is given for m_{H^0}=125 GeV. See also
                                                                                                                                 NODE=S126STT;LINKAGE=A
     CHATRCHYAN 14AJ.
   <sup>2</sup>AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations with up to 10.0 fb<sup>-1</sup> and 9.7 fb<sup>-1</sup>, respectively, of p\overline{p} collisions at E_{\rm cm}=1.96 TeV. The
                                                                                                                                 NODE=S126STT;LINKAGE=AT
     quoted signal strength is given for m_{\mbox{$H^0$}}=125 GeV.
   ^3 AAD 12AI obtain results based on 4.7 fb^{-1} of pp collisions at E_{\rm cm}=7 TeV. The quoted signal strengths are given in their Fig. 10 for m_{\mbox{\it H}^0}=126 GeV. See also Fig. 13
                                                                                                                                 NODE=S126STT;LINKAGE=AA
   4 AALTONEN 13L combine all CDF results with 9.45–10.0 fb<sup>-1</sup> of p\bar{p} collisions at E_{\rm cm}
                                                                                                                                 NODE=S126STT;LINKAGE=LL
     = 1.96 TeV. The quoted signal strength is given for m_{H^0}=125 GeV.
   ^5 ABAZOV 13L combine all D0 results with up to 9.7 fb^{-1} of p\overline{p} collisions at E_{\rm cm}=1.96 TeV. The quoted signal strength is given for m_{\slashed{H^0}}=125 GeV.
                                                                                                                                 NODE=S126STT;LINKAGE=AB
```

 6 CHATRCHYAN 12N obtain results based on 4.9 fb $^{-1}$ of pp collisions at $E_{
m cm}{=}7$ TeV

See also CHATRCHYAN 13Y .

and 5.1 fb $^{-1}$ at $E_{\rm cm}=$ 8 TeV. The quoted signal strength is given for $m_{H^0}=$ 125.5 GeV.

$Z\gamma$ Final State

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<11	95				$pp \rightarrow H^0 X$, 7, 8 TeV
< 9.5	95	² CHATRCHYAN	13 _{BK}	CMS	$pp \rightarrow H^0 X$, 7, 8 TeV

 1 AAD 14J search for $H^0 o Z\gamma o \ell\ell\gamma$ in 4.5 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 20.3 fb $^{-1}$ at $E_{\rm cm}=$ 8 TeV. The quoted signal strength is given for $m_{H^0}=125.5$

 2 GeV. 2 CHATRCHYAN 13BK search for $H^0\to~Z\gamma\to~\ell\ell\gamma$ in 5.0 fb $^{-1}$ of pp collisions at $E_{\rm CM}$ = 7 TeV and 19.6 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. A limit on cross section times branching ratio which corresponds to (4–25) times the expected Standard Model cross section is given in the range $m_{H^0}=120$ –160 GeV at 95% CL. The quoted limit is given for $m_{H^0}=125$ GeV, where 10 is expected for no signal.

NODE=S126SZG NODE=S126SZG

NODE=S126SZG;LINKAGE=A

NODE=S126SZG;LINKAGE=TH

$t\bar{t}H^0$ Production

Signal strengh relative to the Standard Model cross section.

VALUE DOCUMENT ID TECN COMMENT $2.5 \begin{array}{l} +0.9 \\ -0.8 \end{array}$ OUR AVERAGE

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.49
$$^{+6.60}_{-6.28}$$
 3 AALTONEN 13L CDF $p\overline{p}$, 1.96 TeV <5.8 95 4 CHATRCHYAN 13X CMS $pp \rightarrow H^0 t\overline{t} X$

¹ AAD 15 search for $t\bar{t}H^0$ production with H^0 decaying to $\gamma\gamma$ in 4.5 fb⁻¹ of pp collisions at $E_{\rm cm}=7$ TeV and 20.3 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The quoted result on the signal strength is equivalent to an upper limit of 6.7 at 95% CL and is given for $m_{H^0}=125.4$ GeV.

 2 KHACHATRYAN 14H search for $t\bar{t}H^0$ production with H^0 decaying to $b\bar{b},~\tau\tau,~\gamma\gamma,~WW^*$, and ZZ^* , in 5.1 fb $^{-1}$ of pp collisions at $E_{\rm cm}=7$ TeV and 19.7 fb $^{-1}$ at $E_{\rm cm}=8$ TeV. The quoted signal strength is given for $m_{H^0}=125.6$ GeV.

 3 AALTONEN 13L combine all CDF results with 9.45–10.0 fb $^{-1}$ of $p \overline{p}$ collisions at $E_{
m cm}$ = 1.96 TeV. The quoted signal strength is given for $m_{H^0}=125~{\rm GeV}.$

⁴ CHATRCHYAN 13X search for H^0 $t\bar{t}$ production followed by $H^0 \to b\bar{b}$, one top decaying to $\ell\nu$ and the other to either $\ell\nu$ or $q\overline{q}$ in 5.0 fb⁻¹ and 5.1 fb⁻¹ of pp collisions at $E_{\mathrm{cm}}=7$ and 8 TeV. A limit on cross section times branching ratio which corresponds to (4.0–8.6) times the expected Standard Model cross section is given for $m_{H^0}=110$ –140 GeV at 95% CL. The quoted limit is given for $m_{H^0}=125$ GeV, where 5.2 is expected for no signal.

NODE=S126STH NODE=S126STH NODE=S126STH

OCCUR=2

NODE=S126STH:LINKAGE=B

NODE=S126STH;LINKAGE=A

NODE=S126STH;LINKAGE=LL

NODE=S126STH;LINKAGE=TY

H⁰ REFERENCES

			••		
CHATRCHYAN CHATRCHYAN CHATRCHYAN CHATRCHYAN KHACHATRY KHACHATRY AAD AALTONEN ABAZOV CHATRCHYAN AAD AALTONEN AALTONEN	14AS 14BC 14J 14O 14W 14F 14AA 14AI 14B 14B 11AB 13BK 13L 13BK 13L 12AI 12DA 12QA 12CA 12CA 12CA 12CA 12CA 12CA 12CA 12C	PRL 113 161802 PR D89 092007 PR D89 012003 NATP 10 557 EPJ C74 2980 JHEP 1401 096 JHEP 1405 104 PL B736 64 JHEP 1409 087 EPJ C74 3076 PL B726 120 PL B726 120 PL B726 88 PR D88 052014 PR D88 052011 PL B726 587 PRL 110 081803 JHEP 1305 145 JHEP 1306 081 PL B716 1 SCI 338 1576 PRL 109 111804 PRL 109 111805 PRL 109 111805 PRL 109 111805 PRL 109 111805 PRL 109 171804 PRL 109 171804 PRL 109 171804 PRL 109 171805 PRL 109 171807		G. Aad et al. S. Chatrchyan et al. V. Khachatryan et al. V. Khachatryan et al. V. Khachatryan et al. T. Aaltonen et al. T. Aaltonen et al. T. Aaltonen et al. S. Chatrchyan et al. S. Chatrchyan et al. T. Aaltonen et al. S. Chatrchyan et al. S. Chatrchyan et al. S. Chatrchyan et al.	(ATLAS Collab.) (CMS Collab.) (CDF COllab.) (CDF Collab.) (CMS Collab.) (CDF Collab.) (CDS Collab.) (CDS Collab.) (COS Collab.) (CMS Collab.)
				•	,

NODE=S126

REFID = 56266 REFID = 56424 REFID = 56424 REFID = 56125 REFID = 56127 REFID = 556126 REFID = 55849 REFID = 55849 REFID = 56033 REFID = 56033 REFID = 56423 REFID = 56423 REFID = 55733 REFID = 55733 REFID = 55733 REFID = 556423 REFID = 556423 REFID = 556183 REFID = 556183 REFID = 556183 REFID = 55192 REFID = 55192 REFID = 55193 REFID = 55193 REFID = 55493 REFID = 54486 REFID = 54247 REFID = 54248 REFID = 54218 REFID = 54218	51457599558888888551585578971271
---	----------------------------------